

**Docosahexaenoic acid and arachidonic acid are fundamental supplements for
induction of neuronal differentiation**

Inna Kan,¹ Eldad Melamed,¹ Daniel Offen¹ and Pnina Green²

¹Laboratory of Neurosciences and ²Laboratory for the Study of Fatty Acids,
Felsenstein Medical Research Center, Beilinson Campus affiliated with Sackler
School of Medicine, Tel Aviv University, Petah Tiqwa 49100, Israel.

Running footline: Induced neuronal differentiation requires DHA and AA

Correspondence: Pnina Green MD, PhD
Department of Internal Medicine B
Rabin Medical Center, Beilinson Campus
Petah Tiqwa 49100, Israel
Phone: 972-3-9376602
Fax: 972-3-9376612
E-mail: pgreen@post.tau.ac.il

Abbreviations: AA, arachidonic acid; DHA, docosahexaenoic acid; FAME, fatty acid
methyl ester; MSC, mesenchymal stem cells

Abstract

Cell replacement therapy is being investigated for the treatment of neurodegenerative disorders. Adult autologous bone-marrow-derived mesenchymal stem cells (MSCs) have been induced to differentiate into neuron-like cells harboring a variety of neuronal markers and transcription factors. Neural tissue characteristically contains high proportions of docosahexaenoic acid (DHA) and arachidonic acid (AA). In the present study, evaluation of the fatty-acid profile of differentiated neuron-like cells revealed a very low level of DHA, similar to that in MSCs but different from typical neurons. Supplementation of the medium with DHA alone resulted in increased levels of DHA but concomitant low levels of AA. However, supplementation of both DHA and AA yielded a fatty acid profile resembling that of neural tissue. It also resulted in enhanced outgrowth of neurite-like processes, hallmarks of neuronal differentiation. These findings demonstrate the essentiality of DHA and AA supplementation in the process of induced neuronal differentiation and have important implications for the development of cell replacement strategies of neural repair.

Key Words: Mesenchymal stem cells, neurite growth, cell replacement therapy, polyunsaturated fatty acids.

Many neurodegenerative disorders are attributed to the degeneration of specific neuronal populations with subsequent functional loss. Although several treatments have been shown to modify the course of the disease, none successfully halted the degeneration. Owing to these factors, cell replacement therapy to replace the degenerated neural cells may serve as a valuable alternative to achieve significant clinical improvement.

Adult autologous bone-marrow-derived mesenchymal stem cells (MSCs) have been extensively studied as candidates for cell replacement therapy of neurodegenerative diseases. Several laboratories, including ours, have shown that MSCs are able to differentiate beyond tissues of mesodermal origin into neuron-like cells harboring a variety of neuronal markers and transcription factors (1-7). However, none of these studies addressed the unique fatty acid composition of the neuronal membranes, namely, the high proportion of the polyunsaturated fatty acids (PUFA) docosahexaenoic acid (DHA, 22:6 n-3) and arachidonic acid (AA, 20:4 n-6) (8).

DHA and AA are elongation-desaturation products of the parent fatty acids, alpha-linolenic acid (LnA) of the omega 3 (n-3) family (18:3n-3) and linoleic acid (LA) of the omega 6 (n-6) family (18:2n-6), respectively. LnA and LA have been identified as essential fatty acids, since they are not synthesized de novo by animals but obtained from plants by diet. They can, however, undergo a series of elongation-desaturation steps in the mammalian organism to form the longer chain derivatives (9). In neural tissue, PUFA are important for multiple aspects of neuronal development and function, including neurite outgrowth (10-14), signal transduction, and membrane fluidity (15-17). Although DHA is found in abundance in neuronal tissue, it cannot be synthesized by neurons and has to be supplied by cerebrovascular endothelium and astrocytes (18).

In the present study, we examined induced differentiation of MSCs in the absence and presence of PUFA supplements.

Materials and Methods

Isolation and culture of human mesenchymal stem cells

This study was approved by the Helsinki Committee of the Israel Ministry of Health and Tel Aviv University. MSCs were isolated as previously described (7) and

cultured in Growth Medium consisting of Dulbecco's modified Eagle's medium (DMEM; Biological Industries, Bet-Haemek, Israel) supplemented with 15% fetal calf serum (FCS; Biological Industries), 2 mM L-glutamine (Biological Industries), 100 µg/ml streptomycin, 100 U/ml penicillin, and 12.5 U/ml nystatin (SPN; Biological Industries). The medium was replaced twice a week.

Neuronal differentiation

Neuronal differentiation was induced by a previously described protocol (7). Briefly, Growth Medium was replaced with Differentiation Medium I consisting of DMEM supplemented with 10% FCS, 2mM glutamine, SPN, 10 ng/ml basic fibroblast growth factor (bFGF; R&D Systems, Minneapolis, MN), 10 ng/ml epidermal growth factor (EGF; R&D systems), and N2 supplement (5µg/ml insulin; 20 nM progesterone; 100 µM putrescine; 30 nM selenium; 100 µg/ml transferrin). Forty-eight hours later, the medium was replaced with Differentiation Medium II containing DMEM supplemented with SPN, 2 mM L-glutamine, N2 supplement, 200 µM butylated hydroxyanisole (BHA; Sigma, St. Louis, MO), 1 mM dibutyryl cyclic AMP (dbcAMP; Sigma), 3-isobutyl-1-methyl-xanthine (IBMX; Sigma) and 1 µM all-*trans*-retinoic acid (RA; Sigma) for 48 hours.

Fatty acid supplementation and analysis

Various concentrations and combinations of DHA and AA, coupled with 1% horse serum and diluted in DMEM and α -tocopherol (40 µM), dissolved in ethanol, were added to Differentiation Medium I.

Following aspiration of the medium, cultures were washed with PBS, and lipids were extracted with hexane (BioLab, Jerusalem, Israel)/isopropanol (Sigma) (3:2, v/v) containing 5 mg/100 ml butylated hydroxytoluene (Sigma) as an antioxidant and 0.5 mg/100 ml heneicosanoic acid (21:0; Sigma), as an internal standard. Fatty acids were converted to fatty acid methyl esters (FAMES) by heating with 14% boron trifluoride (BF₃) in methanol (Sigma) and separated on capillary columns in an HP 5890 Series II GC equipped with a flame ionization detector. Peak areas were integrated and plotted with the aid of the Varian Star Integrator computer package. Individual FAMES were identified by comparing retention times with authentic standards. The amount of individual fatty acid was documented as the weight percentage of the total 25 identified fatty acids (mean \pm SEM).

Immunocytochemistry

Cells were fixed with 0.4% paraformaldehyde and permeabilized with 0.5% Triton X-100 (Sigma). Immunofluorescent double labeling was performed using mouse anti-microtubule associated protein 2 (MAP2; 1:250; Zymed, San Francisco, CA) and goat anti-mouse CyTM2-conjugated AffiniPure (1:50; Jackson, West Grove, PA). Nuclei were counterstained with 4,6-diamidino-2-phenylindole dihydrochloride (DAPI; Sigma).

Neurite measurements

Cells were stained with anti-MAP2 antibody, to reveal the somatodendritic compartment, and DAPI nuclear stain and photographed with Olympus IX70-S8F2 fluorescence microscope. The total neurite length per neuron was determined by measuring the individual neurite lengths with the Image-Pro® Plus software (Media Cybernetics, Silver Spring, MD) and summing them per neuron. At least 20 non-clustered neuron-like cells were analyzed for each treatment group.

Statistics

The statistical significance of differences in mean neurite length between treatment groups was determined by independent-samples t-test.

Results

Induced neuronal differentiation does not affect cellular concentrations of DHA and AA

The concentrations of DHA and AA in the differentiated neuron-like cells were $3.82 \pm 0.17\%$ and $7.17 \pm 0.30\%$, respectively (Fig. 1). These values did not differ significantly from those of undifferentiated MSCs ($3.00 \pm 0.13\%$ and $8.69 \pm 0.63\%$, respectively) (Fig. 1).

DHA supplementation greatly increases DHA concentration in neuron-like cells, but at the expense of decreased AA

Supplementation of DHA at concentrations of 30, 40, 50 and 60 μM resulted in a continuous, dose-dependent increase in cellular DHA levels. However, a considerable concomitant decrease in the concentration of AA was observed (Fig. 2).

Combined DHA and AA supplementation results in a neuron-like cellular fatty acid composition

To prevent the enhanced decrease of AA in neuron-like cells that was triggered by DHA supplementation, we supplemented the differentiation medium with

both AA and DHA. Following the addition of 30, 40, and 50 μM DHA and equal concentrations, respectively, of AA, the DHA levels rose in a dose-dependent manner, while the concentration of AA remained stable (Fig. 3). The combination of 40 μM DHA and 40 μM AA resulted in a final proportion of $8.93\pm 0.24\%$ DHA and $7.93\pm 0.24\%$ AA in neuron-like cells. These values, as well as their ratio of 1.13 ± 0.05 , were deemed to best resemble the concentrations of these fatty acids in neural tissue. Therefore, the combination of 40 μM DHA and 40 μM AA was chosen for further experiments.

Combined DHA and AA supplementation results in enhanced neurite outgrowth

Cultures supplemented with DHA and AA during induced neuronal differentiation showed an increase in the population of neurons with longer total neurite lengths (50–200 μm and higher) and a decrease in the number of neurons with shorter total neurite lengths (0–50 μm) (Fig. 4). In particular, 33% of the neuron-like cells that were differentiated without PUFA supplementation had total neurite length of 0–50 μm and 67% had total neurite length of 50–200 μm and higher. Neuron-like cells that were differentiated with PUFA supplementation had a different frequency distribution: 9% had total neurite length of 0–50 μm and 91% had total neurite length of 50–200 μm and higher. Moreover, following PUFA supplementation, the sum of the total neurite lengths in 21 neuron-like cells increased from 1432 μm to 2514 μm . The observed difference in total neurite length per neuron between the DHA-treated neuron-like cells and the control cells was statistically significant ($p=0.025$, independent-samples t- test).

Discussion

The present study showed that the fatty acid profile of neuron-like cells derived from MSCs following induced differentiation was notably different from the fatty acid profile of neural tissue (19) and resembled the fatty acid profile of the MSCs. By supplementing the differentiation medium with both DHA and AA, we succeeded in attaining a cellular fatty acid profile of neural tissue. Moreover, these neuron-like cells exhibited enhanced neurite outgrowth.

We hereby report that although differentiated bone marrow derived MSCs attain the characteristics of neuronal cells (7), they lack their typical fatty acid profile

(19), the main difference being a lower level of DHA. The importance of adequate supplementation of DHA to the brain has been amply demonstrated (20). However, neurons cannot perform the final step in DHA biosynthesis and depend on its continuous supply. In the present study, the neuronal induction medium contained very small amounts of AA and almost no DHA (data not shown). Supplementation of DHA increased the cellular DHA concentration, but also led to a decrease in the concentration of AA. This effect of DHA supplementation has been observed previously (21). Since both PUFAs are critical to neuronal function, we decided to add DHA and AA to the differentiation medium in an attempt to achieve the typical neuronal fatty acid composition. Following trials with several dose combinations, we found that supplementing the differentiation medium with 40 μM DHA and 40 μM AA resulted in cellular fatty acid profile values resembling those of neural cells from the region of the striatum (19). In addition, the enriched differentiation yielded the inverted DHA/AA ratio ($\text{DHA/AA} > 1$) that characterizes typical neurons.

Cell cultures of neuronal origin have been previously supplemented with DHA, mainly in experiments exploring the role of DHA in neurons (13, 22). Even though increased concentrations of DHA were achieved, the basic proportions of DHA and AA did not change. This is the first time, however, that non-neuronal cells of bone marrow origin (MSCs), induced to differentiate into neurons, attained the cellular fatty acid profile that characterizes typical neurons. It should be mentioned that in the absence of the neuronal induction, MSCs did not attain a neuronal fatty acid profile following PUFA enrichment of the growth medium (data not shown). Thus, both neuronal induction and supplementation with PUFAs, especially DHA, are necessary for MSCs to attain neuronal characteristics.

One of the critical steps in neuronal differentiation is the outgrowth of neuronal processes (axons and dendrites), which establish the neuron's structural and functional polarity (23, 24). It has previously been demonstrated that in the course of neuronal induction, human MSCs develop cellular features of neuritogenesis and synaptogenesis similar to those observed in immature neurons (25). In the present study, the addition of 40 μM DHA and 40 μM AA to the induction differentiation medium enhanced neurite growth of neuron-like cells. Several studies have demonstrated both an increase in the population of neurons with longer neurites and a higher number of branches following DHA and AA supplementation, indicating a possible role of these PUFAs in promoting neuronal differentiation (10-14). In these

experiments, AA and DHA in concentrations of 1.5–60 μM significantly increased neurite outgrowth in several cell types of neuronal origin. The novelty of our study lies in the fact that the same effect was observed with cells of mesenchymal origin engineered to become neurons. The molecular mechanisms underlying the contribution of DHA and AA to neurite growth are not completely understood, and several possibilities have been raised. These include stimulation of the phospholipid synthesis required for neurite elongation and membrane expansion and modulation of the signal transduction pathways involved in neurite outgrowth. Clearly, further studies are required to establish the exact mechanism(s) of PUFA-induced neurite growth in MSCs undergoing induced neuronal differentiation in vitro.

In conclusion, we demonstrated the need for both DHA and AA supplementation to MSCs in the process of induced neuronal differentiation. The membrane composition of the differentiated cells is an important feature that influences their integration, synaptic formation, and function. Therefore, the attainment of a neuronal fatty acid profile during induced neuronal differentiation is an essential step in the induction of stem cells into functional neurons.

Acknowledgements: We thank Aviva Kluska, Dr. Yossef Levy, Dr. Shlomo Bulvik and Beilinson Editorial Board for their assistance.

This work was supported in part by the Public Committee for the Designation of Estate Funds, the Ministry of Justice, Israel (P.G., D.O.), by the National Parkinson Disease Foundation, U.S.A. (D.O., E.M.) and by the Norma and Alan Aufzein Chair of Research of Parkinson's Disease (E.M.).

References

1. Schwarz, E.J., G.M. Alexander, D.J. Prockop, and S.A. Azizi. 1999. Multipotential marrow stromal cells transduced to produce L-DOPA: Engraftment in a rat model of Parkinson disease. *Hum. Gene Ther.* **10**:2539-2549.
2. Schwarz, E.J., R.L. Reger, G.M. Alexander, R. Class, S.A. Azizi, and D.J. Prockop. 2001. Rat marrow stromal cells rapidly transduced with a self-inactivating retrovirus synthesize L-DOPA in vitro. *Gene* **8**:1214-1223.
3. Woodbury, D., K. Reynolds, and I.B. Black. 2002. Adult bone marrow stromal stem cells express germline, ectodermal, endodermal, and mesodermal genes prior to neurogenesis. *J. Neurosci. Res.* **69**:908-917.
4. Hermann, A, R. Gastl, S. Liebau, M.O. Popa, J. Fiedler, B.O. Boehm, M. Maisel, H. Lerche, J. Schwarz, R. Brenner, and A. Storch. 2004. Efficient generation of neural stem cell-like cells from adult human bone marrow stromal cells. *J. Cell Sci.* **117**:4411-4422.
5. Dezawa, M., H. Kanno, M. Hoshino, H. Cho, N. Matsumoto, Y. Itokazu, N. Tajima, H. Yamada, H. Sawada, H. Ishikawa, T. Mimura, M. Kitada, Y. Suzuki, and C. Ide. 2004. Specific induction of neuronal cells from bone marrow stromal cells and application for autologous transplantation. *J. Clin. Invest.* **113**:1701-1710.
6. Levy, Y.S., D. Merims, H. Panet, Y. Barhum, E. Melamed, and D. Offen. 2003. Induction of neuron-specific enolase promoter and neuronal markers in differentiated mouse bone marrow stromal cells. *J. Mol. Neurosci.* **21**:121-132.
7. Blondheim, N.R., Y.S. Levy, T. Ben-Zur, A. Burshtein, T. Cherlow, I. Kan, R. Barzilai, M. Bahat-Stromza, Y. Barhum, S. Bulvik, E. Melamed, and D. Offen. 2006. Human mesenchymal stem cells express neural genes, suggesting a neural predisposition. *Stem Cells Dev.* **15**:141-164.
8. Innis, S.M. 1991. Essential fatty acids in growth and development. *Prog. Lipid Res.* **30**:39-103.
9. Sprecher, H. 2000. Metabolism of highly unsaturated n-3 and n-6 fatty acids. *Biochim. Biophys. Acta* **1486**:219-231.

10. Dehaut, F., I. Bertrand, T. Miltaud, A. Pouplard-Barthelaix, and M. Maingault. 1993. N-6 polyunsaturated fatty acids increase the neurite length of PC12 cells and embryonic chick motoneurons. *Neurosci. Lett.* **161**:133-136.
11. Okuda, S., H. Saito, and H. Katsuki. 1994. Arachidonic acid: Toxic and trophic effects on cultured hippocampal neurons. *Neuroscience* **63**:691-699.
12. Williams, E.J., F.S. Walsh, and P. Doherty. 1994. The production of arachidonic acid can account for calcium channel activation in the second messenger pathway underlying neurite outgrowth stimulated by NCAM, N-Cadherin, and L1. *J. Neurochem.* **62**:1231-1234.
13. Calderon, F., and H.Y. Kim. 2004. Docosahexaenoic acid promotes neurite growth in hippocampal neurons. *J. Neurochem.* **90**:979-988.
14. Marszalek, J.R., H.F. Lodish. 2005. Docosahexaenoic acid, fatty acid-interacting proteins, and neuronal function: Breastmilk and fish are good for you. *Annu. Rev. Cell Dev. Biol.* **21**:633-657.
15. Jones, C.R., T. Arai, and S.I. Rapoport. 1997. Evidence for the involvement of docosahexaenoic acid in cholinergic stimulated signal transduction at the synapse. *Neurochem. Res.* **22**:663-670.
16. Chalon, S., S. Delion-Vancassel, C. Belzung, D. Guilloteau, A.M. Leguisquet, J.C. Besnard, and G. Durand. 1998. Dietary fish oil affects monoaminergic neurotransmission and behavior in rats. *J. Nutr.* **128**:2512-2519.
17. Bazan, N.G. 2003. Synaptic lipid signaling: significance of polyunsaturated fatty acids and platelet-activating factor. *J. Lipid Res.* **44**:2221-2233.
18. Moore, S.A. 2001. Polyunsaturated fatty acid synthesis and release by brain-derived cells in vitro. *J. Mol. Neurosci.* **16**:195-200.
19. Green, P., I. Gispan-Herman, and G. Yadid. 2005. Increased arachidonic acid concentration in the brain of Flinders Sensitive Line rats, an animal model of depression. *J. Lipid Res.* **46**:1093-1096.
20. McNamara, R.K., and S.E. Carlson. 2006. Role of omega-3 fatty acids in brain development and function: Potential implications for the pathogenesis and prevention of psychopathology. *Prostaglandins Leukotr. Essent. Fatty Acids* **75**:329-349.
21. Wainwright, P.E., H.C. Xing, G.R. Ward, Y.S. Huang, E. Bobik, N. Auestad, and M. Montalto. 1999. Water maze performance is unaffected in artificially

- reared rats fed diets supplemented with arachidonic acid and docosahexaenoic acid. *J. Nutr.* **129**:1079-1089.
22. Kim, H.Y., M. Akbar, A. Lau, and L. Edsall. .2000. Inhibition of neuronal apoptosis by docosahexaenoic acid (22:6n-3). Role of phosphatidyl-serine in antiapoptotic effect. *J. Biol. Chem.* **275**:35215-35223.
 23. Cline, H.T. 2001. Dendritic arbor development and synaptogenesis. *Curr. Opin. Neurobiol.* **11**:118-126.
 24. Poirazi, P., and B.W. Mel. 2001. Impact of active dendrites and structural plasticity on the memory capacity of neural tissue. *Neuron* **29**:779-796.
 25. Wensch, S., K. Trinkaus, A. Hild, D. Hose, C. Heiss, and V. Alt. 2006. Immunochemical, ultrastructural and electrophysiological investigations of bone-derived stem cells in the course of neuronal differentiation. *Bone* **38**: 911–921.

Footnote

This work was performed in partial fulfillment of the requirements for a Ph.D. degree of I.K.



Figure Legends

Figure 1. DHA and AA concentrations in MSCs before and after induced neuronal differentiation. The fatty acid composition of the cells was determined by gas chromatography of fatty acid methyl esters, as described in Materials and Methods. DHA, docosahexaenoic acid; AA, arachidonic acid; MSCs, mesenchymal stem cells before induced neuronal differentiation (n=18); Diff., MSCs following induction of differentiation (n=33). Values are expressed as weight percentage of the total identified fatty acids, means \pm SEM.

Figure 2: DHA and AA concentrations following supplementation with DHA. Neuronal differentiation was induced either without DHA (0 μ M) or with supplementation of increasing concentrations of DHA (30, 40, 50 and 60 μ M), as described in Materials and Methods. The DHA and AA composition of the cells was determined by gas chromatography of fatty acid methyl esters, as described in Materials and Methods. DHA, docosahexaenoic acid; AA, arachidonic acid. Values are weight percentages of the total identified fatty acids, means \pm SEM (n=6).

Figure 3: DHA and AA concentrations following supplementation with both DHA and AA. Differentiation was induced either without PUFAs (Diff.) or with supplementation of different concentrations (Diff. + PUFA) of DHA and AA, as specified in Materials and Methods. Diff., MSCs following induced neuronal differentiation; PUFA, polyunsaturated fatty acids; DHA, docosahexaenoic acid; AA, arachidonic acid. Values are weight percentages of the total identified fatty acids, means \pm SEM (n=5).

Figure 4. Effect of DHA and AA supplementation on neurite length of differentiated cells. Neurite length of differentiated cells was determined in the absence (Diff.) and presence of 40 μ M DHA and 40 μ M AA (Diff. + PUFA), as described in Materials and Methods. Diff., MSCs following induced neuronal differentiation; PUFA, polyunsaturated fatty acids

Fig. 1





